

REMARKS

The examiner's stated basis for restriction is that Nagase *et al.*, *DNA Research* 3: 43-53 (1996) ("Nagase II"), "teaches a KIAA0172 gene," which the examiner asserts to be the "technical feature linking groups I – VII" (action at page 3, lines 1 – 3). With this perspective, the examiner concludes that the "technical feature" in question "does not define a contribution over the art" (*id.*, lines 5 – 6).

Applicants would point out, however, that Nagase I is supplemental to another, contemporaneous article by Nagase *et al.*, *DNA Research* 3: 17-24 (1996) ("Nagase I"), which is Exhibit A to this response. Nagase I discloses a cDNA clone, designated "KIAA0172," said to encode a polypeptide of 1307 amino acids (see Table 3 of Nagase I).

The polypeptide thus disclosed actually does not exist in any intact cell, however; this, because Nagase I makes an erroneous assignment of the first codon. Instead, for a correct assignment of the first codon the mRNA cap site had to be determined, via extensive and elaborate experimental investigation, by the present inventors.

Accordingly, the present application confirms the absence from cells of a polypeptide as taught by Nagase I. Likewise incorrect is the GenBank sequence, D79994, that corresponds to the Nagase I disclosure and that the authors deposited. A copy of the D79994 sequence is Exhibit B to this response.

It is apparent, therefore, that the present application is the first to disclose a correct sequence for "a KIAA0172 gene." That correct sequence is applicants' SEQ ID NO: 1, which has 1194 amino acids.

This 1194 amino-acid sequence differs from the Nagase I sequence, which has 1367 amino acids. Furthermore, neither Nagase I nor Nagase II discloses a function for the protein encoded by KIAA0172. Indeed, the protein posited by Nagase I differs so markedly from the protein having the sequence of SEQ ID NO:1 that one could not reasonably predict that the former has the same function as the present application describes for the latter.

In light of the foregoing, it is evident that the examiner's stated basis for restriction is factually untenable. Accordingly, what he has identified as the technical feature linking the invention of groups I – VIII in fact does define a contribution over the prior art represented by Nagase I/II. The respective claim sets of Groups I – VIII should not be divided, therefore, but rather should be examined together.

Respectfully submitted,

Date 10 July 2006

By S. A. Bent

FOLEY & LARDNER LLP
Customer Number: 22428
Telephone: (202) 672-5404
Facsimile: (202) 672-5399

Stephen A. Bent
Attorney for Applicant
Registration No. 29,768

Prediction of the Coding Sequences of Unidentified Human Genes. V. The Coding Sequences of 40 New Genes (KIAA0161-KIAA0200) Deduced by Analysis of cDNA Clones from Human Cell Line KG-1

Takahiro NAGASE, Naohiko SEKI, Ken-ichi ISHIKAWA, Ayako TANAKA, and Nobuo NOMURA*
Kazusa DNA Research Institute, 1532-3 Yanauchino, Kisarazu, Chiba 292, Japan

(Received 1 February 1996; revised 21 February 1996)

Abstract

As part of our continuing efforts to accumulate information on the coding region of unidentified human genes, we newly determined the sequences of 40 cDNA clones of human cell line KG-1 which correspond to relatively long and nearly full-length transcripts, and predicted the coding sequences of the corresponding genes, named KIAA0161 to 0200. The average size of the cDNA clones analyzed was approximately 5.0 kb. A computer search of the sequences in public databases indicated that the sequences of 20 genes were unrelated to any reported genes, while the remaining 20 genes carried sequences which show some similarities to known genes. Among the genes in the latter category, KIAA0167 contained a Zn-finger motif with significant structural similarity to that of the yeast transcription factor *GCSI*, and KIAA0189 was classified into the RhoGAP gene family. Stretches of typical CAG (Gln) repeats, which were often correlated with genetic disorders, were found in KIAA0181 and KIAA0192. Another novel repeat composed of alternating Arg and Glu was identified in KIAA0182. Northern hybridization analysis demonstrated that 10 genes are expressed in a cell- or tissue-specific manner.

Key words: full-length cDNA sequence; CAG repeat; transcriptional factor; RhoGAP gene family; myeloid cell line KG-1

1. Introduction

In this series of projects involving the accumulation of information on the coding sequences of unidentified human genes, we have been analyzing nearly full-length cDNA clones which were isolated from human immature myeloid cell line KG-1.¹ We already reported the sequences of 160 new cDNA clones and predicted the coding regions of the corresponding genes.¹⁻⁴ The average size of these cDNA clones, except for 20 clones (KIAA0101 to 0120), was approximately 4.0 kb. Each clone contained a distinct open reading frame (ORF) in the 5'-moiety, and their average size was roughly 1.7 kb, indicating that most of the clones carried relatively long 3'-untranslated regions (3'-UTRs). Although vast amounts of expressed sequence tags (ESTs) obtained by single-run sequencing of cDNA libraries have been accumulated for comprehensive understanding of expression profiles, our preliminary analysis indicated that most of ESTs (GenBank release 92.0, Dec. 1995) fell in the region about 2 kb from the poly(A)-tail of our cDNA sequences. This is probably due to the fact that the cDNA libraries prepared by conventional methods contain a fairly large amount of small clones derived from

truncated transcripts. On the basis of computer analysis, biological function has been predicted for at least 40% of the genes that we reported, from which the functional significance of 20 genes is under investigation in collaboration with other laboratories. We keep on sequencing the new cDNA clones, and in this paper, we report the coding sequences of 40 additional genes and their sequence features as well as expression profiles.

2. Materials and Methods

The source of cDNA libraries and methods used for selection of cDNA clones, Northern hybridization, sequence analysis, computer analysis of sequences and chromosomal mapping of cDNA clones were described previously.¹

3. Results and Discussion

3.1. Sequence features of analysed cDNA clones

The cDNA clones carrying inserts longer than 2 kb were randomly selected from the libraries constructed from the middle-sized cDNA class, and both the terminal sequences were analysed to select unidentified clones with poly(A) tails.¹ The clones carrying inserts which were more than 90% of the length of the corresponding transcripts were further selected by Northern hybridization,

Communicated by Mituru Takanami

* To whom correspondence should be addressed. Tel. +81-438-52-3930, Fax. +81-438-52-3931




Fig. 1. Physical maps of the 40 cDNA clones analyzed. The horizontal scale represents the cDNA length in kb, and gene numbers are given on the left. Open reading frames (ORFs) within coding regions, untranslated regions and repetitive sequences are indicated by solid, open and dotted boxes, respectively. The positions of the first ATG codon in each ORF are represented by triangles. The names of repetitive sequences are described above the dotted boxes. The solid bars show the positions of the triplet and other repeats listed in Fig. 2. The nucleotide sequence data reported in this paper were deposited in the GSDB, DDBJ, EMBL and NCBI nucleotide sequence databases under the accession numbers shown in Table 3.

Table 1. cDNA clones with similarities to PIR and GenBank/EMBL database files.

Gene no. (KIAA)	Database files	Accession no. ^{a)}	Identity (%)	Overlap ^{b)} (amino acid residues)
0162	emb-5 protein	S35241	31.5	1525
0164	DNA-binding protein 5 (H)	S26650	38.4	159
0165	cut1 protein (Sp)	A35694	36.0	236
0167	KIAA0041 (H)	D26069 ^{c)}	46.0	126
	KIAA0050 (H)	D30758 ^{c)}	35.5	166
0170	calphotin (D)	A47283	17.8	841
0171	hypothetical protein L8167.6 (Sc)	S48557	30.8	214
0172	ankyrin 3, long form (H)	A55575	34.5	139
0173	tubulin-tyrosine ligase (P)	A45443	30.4	214
0175	protein kinase p69Eg3 (X)	S52244	62.1	655
0177	poly(ADP-ribose)synthase (C)	JH0581	29.0	303
0178	SMC1 protein (Sc)	A49464	28.2	1241
0179	hypothetical protein D4478 (Sc)	S48776	33.8	130
0185	hypothetical protein YM9959.11C (Sc)	S57596	26.0	1087
0188	SMP2 protein (Sc)	S30911	47.9	240
0189	regulator protein p122 - RhoGAP (R)	S54293	40.1	1039
0190	Ubiquitin-specific proteinase UBP3 (Sc)	B44450	23.9	368
0192	Mopa box protein (M)	A26892	94.6	129
0198	finger protein (clone XlcOF7.1) (X)	S06546	29.0	214
0199	hydroxymethylglutaryl-CoA reductase (Su)	A31898	28.0	168
0200	neurogenic locus mam protein (D)	A36391	18.5	243

^{a)} PIR database files are shown except for KIAA0167. ^{b)} The size of regions which show similarities. ^{c)} GenBank/EMBL database files.

C, chicken; Ce, *Caenorhabditis elegans*; D, *Drosophila melanogaster*; H, human; M, mouse; P, pig; R, rat; Sc, *Saccharomyces cerevisiae*; Sp, *Schizosaccharomyces pombe*; Su, sea urchin; X, *Xenopus laevis*.

Table 2. cDNA clones with regions that matched motifs in the PROSITE database.

Motifs	Description	Gene number (KIAA)	References
ZINC FINGER C3HC4	Zinc finger, C3HC4 type	0161	13
ATP GTPA	ATP/GTP-binding site motif A (P-loop)	0167, 0178, 0187	14
PROTEIN KINASE ST	Protein kinases	0175	15
ABC TRANSPORTER	ABC transporters family	0178	16
CRYSTALLIN BETAGAMMA	Crystallin β and γ 'Greek Key' motif	0184	17
UCH 22	Ubiquitin carboxyl-terminal hydrolases family 2	0190	18
ZINC FINGER C2H2	Zinc finger, C2H2 type	0192	19
TOPOISOMERASE II	DNA topoisomerase II	0192	20
HIS ACID PHOSPHAT	Histidine acid phosphatases	0197	21
G BETA REPEATS	β -transducin family Trp-Asp repeats	0199	22

and their sequences were determined.¹ By ORF analysis, each clone was found to contain a distinct ORF. The ORFs and the first ATG codon are shown in Fig. 1 by solid boxes and open triangles, respectively. In-frame termination codons upstream of the first ATG codon were identified in 20 clones, suggesting that at least 50% of the clones analyzed harbor the complete coding region.

The results of computer analysis with the GCG software package⁵ are shown in Tables 1 and 2 and also in the figure in the Supplement section. Sequence features

noted are summarized as follows.

1. Sequences of 20 genes were unrelated to any reported genes, except for EST sequences in the database files. The remaining 20 genes carried sequences with some similarities to known genes (Table 1). Among the genes in the latter category, KIAA0167 contained a Zn-finger motif with significant structural similarity to that of the yeast transcription factor *GCS1*⁶ (indicated by solid line in Fig. 2A). In ad-



Fig. 2. Sequence comparison in the region spanning the Zn-finger motif (solid line) and ankyrin repeat-like sequence (dotted line) in the KIAA0167 gene family and the *GCSI* gene (A) and that spanning the domain sequence of the RhoGAP and three KIAA clones (B). Identical and similar amino acids are indicated by black and grey background, respectively. Numerals indicate the number of amino acid residues from the start codon.

dition, an ankyrin repeat-like sequence⁷ was identified in the adjacent region (indicated by dotted line in Fig. 2A). We noted that two previously reported cDNAs, KIAA0041 and KIAA0050, also carry similar sequences in the corresponding regions. Particular conservation of the Zn-finger motif and ankyrin repeat-like sequence suggests that these genes constitute a novel gene family related to the *GCS1* gene. We tentatively named it the KIAA0167 family. As another gene with similarity to known genes, we noted that KIAA0189 contains the domain sequence of the RhoGAP gene family⁸ (GAP: GTPase-activating proteins) (Fig. 2B). Since the sequence conservation between the two genes is quite high, it is likely that KIAA0189 is a member of this family. We also noted that 2 previously reported genes,

KIAA0013 and KIAA0053, carry sequences showing weak similarity to the GAP domain (Fig. 2B).

2. Protein motifs that matched those in the PROSITE motif database were found in 11 genes (Table 2).
3. Significant transmembrane domains were identified in 13 genes, 5 of which harbored multiple hydrophobic regions, as judged by the methods of Engelman et al.⁹ and of Kyte and Doolittle.¹⁰
4. Two genes harbored stretches of CAG (Gln) repeats, which were often correlated with genetic disorders:¹¹ CAG occurred 23 times within a 42 triplet stretch in KIAA0181 and 62 times within a 100 triplet stretch in KIAA0192 (Fig. 3A and B).
5. Another novel repeat composed of alternating Arg

Table 3 Summary of cDNA sequence data and expression patterns of cloned genes in human tissues and cell lines.

Gene number (KIAA)	Total length of cDNA (bp) ^{a)}	Amino acid residues	Expression ^{b)}																Chromosomal Accession ^{c)}				
			KC-1	HeLa	He	Br	Pl	Lu	Li	Sk.m	Ki	Pa	Sp	Th	Pr	Tc	Ov	Sm.1	Co	Pc	b	location	number
0161 ^{d)}	5,559	292	+	+	-	+	+	-	+	+	+	+	+	-	±	+	±	-	±	±	±	2	D79883
0162 ^{e)}	5,876	1,726	+	+	+	+	+	+	++	+	+	+	+	+	+	+	+	+	+	+	+	17	D79884
0163 ^{d)}	4,436	550	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	22	D79885
0164 ^{e)}	5,538	920	+	+	+	+	+	+	++	+	+	+	+	+	+	+	+	+	+	+	+	6	D79886
0165 ^{d,e)}	6,652	1,795	+	+	+	-	-	-	-	-	-	-	-	-	++	-	+	+	+	-	-	8	D79887
0166 ^{d)}	6,942	2,209	+	+	+	±	±	±	±	±	±	±	±	±	++	±	+	+	+	+	+	17	D79888
0167 ^{e,f)}	3,950	836	+	-	-	++	-	±	±	+	-	-	+	+	-	-	±	-	+	+	+	12	D79889
0168	5,426	326	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	20	D79890
0169 ^{d)}	5,656	1,745	+	+	+	+	+	+	+	+	+	+	+	+	++	+	+	+	+	+	+	9	D79891
0170 ^{e)}	6,940	2,089	+	+	+	+	+	+	++	+	+	+	+	+	++	+	+	+	+	+	+	6	D79892
0171 ^{e)}	3,336	625	+	+	+	+	+	+	++	+	+	+	+	+	+	+	+	+	+	+	+	5	D79893
0172 ^{e)}	4,792	1,307	+	+	+	±	±	±	+	+	+	+	+	+	+	+	+	+	+	+	+	9,20	D79894
0173 ^{e,f)}	4,831	1,199	+	+	+	±	±	±	+	+	+	+	+	+	++	+	+	+	+	+	+	2	D79895
0174	2,348	361	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	19	D79896
0175 ^{d,e,f)}	2,470	651	+	+	+	+	+	-	-	-	-	-	-	-	+	±	±	+	+	+	-	9	D79897
0176	3,635	263	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	17	D79898
0177 ^{d,e)}	5,083	1,631	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	13,22	D79899
0178 ^{e,f)}	5,803	1,225	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	X	D80000
0179 ^{e)}	4,994	762	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	21	D80001
0180	5,009	343	+	+	+	+	+	+	+	+	+	+	+	+	++	+	+	+	+	+	+	20	D80002
0181	6,504	2,005	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	20	D80003
0182	7,133	1,157	+	+	+	+	+	+	+	+	+	+	+	+	-	+	+	+	+	+	+	16	D80004
0183	4,905	1,062	++	++	++	+	+	++	++	++	++	++	++	++	+	++	+	+	+	+	+	9	D80005
0184 ^{f)}	4,639	863	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	2	D80006
0185 ^{e)}	5,823	1,884	+	+	±	±	±	±	±	±	±	±	±	±	++	+	+	+	+	+	+	10	D80007
0186	3,248	196	+	+	-	-	-	-	-	-	-	-	-	-	+	-	±	±	-	-	-	20	D80008
0187 ^{f)}	4,181	1,282	+	+	++	+	+	+	++	+	+	+	+	+	+	+	+	+	+	+	+	10	D80009
0188 ^{e)}	5,307	899	+	+	+	+	+	+	++	+	+	+	+	±	+	+	+	+	+	+	+	2	D80010
0189 ^{e)}	4,824	1,023	+	-	+	-	+	+	+	+	+	+	±	+	+	+	+	+	+	-	-	X	D80011
0190 ^{e,f)}	3,280	839	+	+	+	+	+	+	+	+	+	+	+	+	++	+	+	+	+	+	+	16,22	D80012
0191 ^{f)}	5,203	1,516	+	+	+	±	±	+	+	+	+	+	+	+	++	-	+	±	±	±	±	1	D83776
0192 ^{d,e,f)}	6,604	2,124	+	+	+	±	±	+	±	+	±	+	+	±	±	+	±	±	±	±	±	X	D83783

Table 3. Continued.

Gene number (KIAA)	Total length of cDNA (bp) ^{a)}	Amino acid residues	Expression ^{b)}													Chromosomal location	Accession number ^{c)}
			KG-1	HeLa	He	Br	Pl	Lu	Li	Sk.m	Ki	Pa	Sp	Th	Pr		
0193	5,076	346	+	+	+	+	+	+	+	+	+	+	+	+	+	7	D83777
0194	5,245	1,435	+	+	+	+	+	+	+	+	+	+	+	+	+	4.5	D83778
0195 ^{d)}	5,022	1,356	+	+	+	+	+	+	+	+	+	+	+	+	+	17	D83779
0196 ^{d)}	4,103	1,139	+	+	+	+	+	+	+	+	+	+	+	+	+	8	D83780
0197 ^{d,e)}	4,814	1,314	+	+	+	+	+	+	+	+	+	+	+	+	+	11	D83781
0198 ^{d,e)}	5,638	562	+	+	+	+	+	+	+	+	+	+	+	+	+	20	D83784
0199 ^{d,e,f)}	3,976	1,277	+	+	+	+	+	+	+	+	+	+	+	+	+	3	D83782
0200 ^{d)}	5,713	1,016	+	+	+	+	+	+	+	+	+	+	+	+	+	5	D83785

n.d., not determined; He, heart; Br, brain; Pl, placenta; Lu, lung; Li, liver; Sk.m, skeletal muscle; Ki, kidney; Pa, pancreas; Sp, spleen; Th, thymus; Pr, prostate; Te, testis; Ov, ovary; Sm.i, small intestine; Co, colon; Pe.b, peripheral blood leukocytes. ^{a)} Values excluding poly(A) sequences. ^{b)} Expression of mRNA in indicated cells and human tissues (Clontech, USA) was examined by northern hybridization, and the strength of the positive signals are indicated (±, +, ++, +++). ^{c)} Accession number of GSDB, DDBJ, EMBL and NCBI nucleotide sequence databases. ^{d)} Putative transmembrane domains were contained (see Supplemental pages). ^{e)} Similarities to known genes were identified (see Table 1 and Supplemental pages). ^{f)} Protein motifs were recognized (see Table 2 and Supplemental pages).

remaining 10 genes apparently showed different expression profiles among the cells and tissues examined. Although the spectra were different for each gene, most of them seemed to belong to the class of genes of which the expression was specifically suppressed in certain tissues. In contrast, it was significant that 4 genes, KIAA0165, 0167, 0175 and 0186 were only expressed in a few specific tissues.

As seen in the previously reported genes (Fig. 2 in ref. 2, 3), multiple but discrete bands were recognized in 30 clones, possibly due to either alternative splicing, alternative termination, or initiation of transcription at different sites.

The chromosomal location of these genes have been determined using a panel of human-rodent hybrid cell lines (see Table 3).

Acknowledgments: This project was supported by grants from the Kazusa DNA Research Institute Foundation. We thank Dr. M. Takanami for his support of this work.

References

1. Nomura, N., Miyajima, N., Sazuka, T. et al. 1994, Prediction of the coding sequences of unidentified human genes. I. The coding sequences of 40 new genes (KIAA0001-KIAA0040) deduced by analysis of randomly sampled cDNA clones from human immature myeloid cell line KG-1, *DNA Res.*, **1**, 27-35, Supplement; **1**, 47-56.
2. Nomura, N., Nagase, T., Miyajima, N. et al. 1994, Prediction of the coding sequences of unidentified human genes. II. The coding sequences of 40 new genes (KIAA0041-KIAA0080) deduced by analysis of cDNA clones from human cell line KG-1, *DNA Res.*, **1**, 223-229, Supplement; **1**, 251-262.
3. Nagase, T., Miyajima, N., Tanaka, A. et al. 1995, Prediction of the coding sequences of unidentified human genes. III. The coding sequences of 40 new genes (KIAA0081-KIAA0120) deduced by analysis of cDNA clones from human cell line KG-1, *DNA Res.*, **2**, 37-43, Supplement; **2**, 51-59.
4. Nagase, T., Seki, N., Tanaka, A. et al. 1995, Prediction of the coding sequences of unidentified human genes. IV. The coding sequences of 40 new genes (KIAA0121-KIAA0160) deduced by analysis of cDNA clones from human cell line KG-1, *DNA Res.*, **2**, 167-174, Supplement; **2**, 199-210.
5. Devereux, J., Haeblerli, P., and Smithies, O. 1984, A comprehensive set of sequence analysis programs for the VAX, *Nucleic Acids Res.*, **12**, 387-395.
6. Ireland, L. S., Johnston, G. C., Drebot, M. A. et al. 1994, A member of a novel family of yeast Zn-finger proteins mediates the transition from stationary phase to cell proliferation, *EMBO J.*, **13**, 3812-3821.
7. Lambert, S., Yu, H., Prchal, J. T. et al. 1990, cDNA sequence for human erythrocyte ankyrin, *Proc. Natl. Acad. Sci. USA.*, **87**, 1730-1734.
8. Homma, Y. and Emori, Y. 1995, A dual functional signal mediator showing RhoGAP and phospholipase C- δ stimulating activities, *EMBO J.*, **14**, 286-291.
9. Engelman, D. M., Steize, T. A., and Goldman, A. 1986, Identifying nonpolar transbilayer helices in amino acid sequences of membrane proteins, *Annu. Rev. Biophys. Biophys. Chem.*, **15**, 321-353.
10. Kyte, J. and Doolittle, R. F. 1982, A simple method for displaying the hydropathic character of a protein, *J. Mol. Biol.*, **157**, 105-132.
11. Mandel, J.-L. 1994, Trinucleotide diseases on the rise, *Nature genetics*, **7**, 453-455.
12. Surowy, C. S., Hoganson, G., Gosink, J., Strunk, K., and Spritz, R. A. 1990, The human RD protein is closely related to nuclear RNA-binding proteins and has been highly conserved, *Gene*, **90**, 299-302.
13. Haupt, Y., Alexander, W. S., Barri, G., Klinken, S. P., and Adams, J. M. 1991, Novel zinc finger gene implicated as *myc* collaborator by retrovirally accelerated lymphomagenesis in *Ep-myc* transgenic mice, *Cell*, **65**, 753-763.
14. Walker, J. E., Saraste, M., Runswick, M. J., and Gay, N. J. 1982, Distantly related sequences in the α - and β -subunits of ATP synthase, myosin, kinases and other ATP-requiring enzymes and a common nucleotide binding fold, *EMBO J.*, **1**, 945-951.
15. Hunter, T. 1991, Protein Kinase Classification, In: Hunter, T., Sefton, B. M. (eds) *Methods in Enzymology*, Vol. 200. Academic Press, New York, pp. 3-37.
16. Higgins, C. F., Hiles, I. D., Salmond, G. P. C. et al. 1986, A family of related ATP-binding subunits coupled to many distinct biological processes in bacteria, *Nature*, **323**, 448-450.
17. Wistow, G. 1993, Lens crystallins: gene recruitment and evolutionary dynamism, *TIBS*, **18**, 301-306.
18. Papa, F. R. and Hochstrasser, M., 1993, The yeast *DOA4* gene encodes a deubiquitinating enzyme related to a product of the human *tre-2* oncogene, *Nature*, **366**, 313-319.
19. Rosenfeld, R. and Margalit, H. 1993, Zinc fingers: conserved properties that can distinguish between spurious and actual DNA-binding motifs, *J. Biomol. Struct. Dyn.*, **11**, 557-570.
20. Wyckoff, E., Natalie, D., Nolan, J. M., Lee, M., and Hsieh, T., 1989, Structure of the *Drosophila* DNA Topoisomerase II gene: Nucleotide sequence and homology among Topoisomerase II, *J. Mol. Biol.*, **205**, 1-13.
21. Van Etten, R. L., Davidson, R., Stevis, P. E., MacArthur, H., and Moore, D. L., 1991, Covalent structure, disulfide bonding, and identification of reactive surface and active site residues of human prostatic acid phosphatase, *J. Biol. Chem.*, **266**, 2313-2319.
22. Gilman, A. G. 1987, G proteins: Transducers of receptor-generated signals, *Annu. Rev. Biochem.*, **56**, 615-649.



All Databases

PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

Search **Nucleotide**for **D79994****Go****Clear****Save**

Limits

Preview/Index

History

Clipboard

Details

Display **Summary**Show **20**

Send to

[About Entrez](#)[Entrez Nucleotide](#)[Help](#) | [FAQ](#)[Entrez Tools](#)[Check sequence
revision history](#)[LinkOut](#)[My NCBI \(Cubby\)](#)[Related resources
BLAST](#)[Reference sequence
project](#)[Search for Genes](#)[Submit to GenBank](#)[Search for full length
cDNAs](#)**All: 1** bacteria: 0 mRNA: 1 RefSeq: 0 Show only records from: [CoreNucleotide \(1\)](#), [EST \(0\)](#), [GSS \(0\)](#). [\[What's this?\]](#)☐ **1: D79994** Reports**Homo sapiens KIAA0172 mRNA, complete cds**
gi|58257638|dbj|D79994.2|[58257638][Links](#)[Disclaimer](#) | [Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)

Jun 21 2006 12:14:26



Nucleotide

My NCBI

[\[Sign In\]](#)
[\[Register\]](#)

PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

Books

Search

Nucleotide

for

Go

Clear

Limits

Preview/Index

History

Clipboard

Details

Display

GenBank

Show

5

Send to

Range: from

begin

to end

☐ Reverse complemented strand

Features: +

Refresh

1: D79994. Reports Homo sapiens KIAA...[gi:58257638]

[Links](#)

[Comment](#) [Features](#) [Sequence](#)

LOCUS D79994 5635 bp mRNA linear PRI 28-JAN-2005
 DEFINITION Homo sapiens KIAA0172 mRNA, complete cds.
 ACCESSION D79994
 VERSION D79994.2 GI:58257638
 KEYWORDS
 SOURCE Homo sapiens (human)
 ORGANISM Homo sapiens
 Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Euteleostomi;
 Mammalia; Eutheria; Euarchontoglires; Primates; Haplorrhini;
 Catarrhini; Hominidae; Homo.
 REFERENCE 1
 AUTHORS Nagase,T., Seki,N., Ishikawa,K., Tanaka,A. and Nomura,N.
 TITLE Prediction of the coding sequences of unidentified human genes. V.
 The coding sequences of 40 new genes (KIAA0161-KIAA0200) deduced by
 analysis of cDNA clones from human cell line KG-1
 JOURNAL DNA Res. 3 (1), 17-24 (1996)
 PUBMED 8724849
 REFERENCE 2
 AUTHORS Chiang,P.W., Wang,S., Smithivas,P., Song,W.J., Ramamoorthy,S.,
 Hillman,J., Puett,S., Van Keuren,M.L., Crombez,E., Kumar,A.,
 Glover,T.W., Miller,D.E., Tsai,C.H., Blackburn,C.C., Chen,X.N.,
 Sun,Z., Cheng,J.F., Korenberg,J.R. and Kurnit,D.M.
 TITLE Identification and analysis of the human and murine putative
 chromatin structure regulator SUPT6H and Supt6h
 JOURNAL Genomics 34 (3), 328-333 (1996)
 PUBMED 8786132
 REFERENCE 3 (bases 1 to 5635)
 AUTHORS Ohara,O., Nagase,T., Kikuno,R. and Nomura,N.
 TITLE Direct Submission
 JOURNAL Submitted (12-DEC-1995) Osamu Ohara, Kazusa DNA Research Institute;
 Kazusa-kamatari 2-6-7, Kisarazu, Chiba, 292-0818, Japan
 (E-mail:cdnainfo@kazusa.or.jp, Tel:81-438-52-3913)
 COMMENT On Jan 27, 2005 this sequence version replaced gi:1136403.
 FEATURES
 source
 Location/Qualifiers
 1..5635
 /organism="Homo sapiens"
 /mol_type="mRNA"
 /db_xref="taxon:9606"
 /clone="ha02512s1"
 /cell_line="KG-1"
 /cell_type="myeloblast"
 /clone_lib="pBlucscriptII SK plus"
 /note="This sequence was obtained by subcloning of the DNA
 fragments derived from two cDNA clones (1-1719 was derived
 from hhl4360 and 1719-5635 was derived from ha2512)."
 gene
 1..5635
 /gene="KIAA0172"

CDS

```
<667..4767
/ gene="KIAA0172"
/ note="Start codon is not identified
similar to ankyrin of Chromatium vinosum."
/ codon_start=1
/ protein_id="AAA11489.2"
/ db_xref="GI:58257639"
/ translation="LLTPFWISHWTQASMAHTTKVNGSASGKAGDILSGDQDKEQKDP
YFVETPYGYQLDLDFLKYVDDIQKGNITKRLNIQKRRKPSVPCPEPRTTSGQQGIWTS
TESLSSNSDDNKQCPNFLIARSQVSTPISKPPPLETSLPFLTIPENRQLPPSPQ
LPKHNHVTKTLMETRRRLERATMQMTPGEFRRPRLASFGGMGTSSLPSSFVSGSN
HNPAAKHQLQNGYQNGDYGSAAPPTSSMGSSIRHSPSSGISTPVTVNSPMHLQH
IREQMAIALKRLKELEEQVRTIPVLQVKISVLQEEKQLVSQLKNQRAASQINVCGR
KRSYSAGNASQLEQLSRARRSGGELYIDYEEEMETVEQSTQRIKEFRQLTADMQALE
QKIQDSSCEASSELRENGECRSVAVGAENMNDIVVYHRGSRCKDAVGTLMVMRNC
GVSVTEAMLGVMTEADKEIELQQQTIEALKEKIYRLEVQLRETHDREMTKLKQELQA
AGSRKKVDKATMAQPLVFSKVVEAVVQTRDQMVGSHMDLVDTCVGTSVETNSVGISQ
PECKNKVVGPELPMNWIVKVERVMHDCAGRSVEMCDKSVSEVSVCTGSNTEESV
NDLTLTKTNLNLKEVRSIGCGDCSVDVTVCSPEKASRGVNTAEVSVQVEAAVMVPR
ADQDTSTDLEQVHQFTNTETATLIESCTNTCLSTLDKQTSTQTVETRTVAVGEGRVK
INSSTKTRSIGVGTLLSGHSGFDRPSAVKTKESGVGQININDNYLVGLKMRITACGP
QLTVGLTASRRSVGVGDDPVGESLENPQPAPLGMMTGDLHYIERIQKLLAEQQTLLA
ENYSELAEAFGEFHSQMGSLSNQLISTLSSINSVMKSASTEELRNPDFQKTSLGKITG
NYLGYTCCKGGLQSGSPLSSQTSQPEQEVGTSEGKPISSLDAPFTQEGTSLSPVNLTD
QIAAGLYACTNNESTLKSIMKKDGNKDSNGAKKNLQFVGINGGYETSSDSSDES
SSSESDDCEDVIEYPLEEEEEDEDEDTRGMAEGHHAVNIEGLKSARVEDEMQVQCEP
EKVEIRERYELSEKMLSACNLLKNTINDPKALTSKDMRFLNTLQHEWFRVSSQKSAI
PAMVGDYIAAFEAISPDVLRVYNLADGNGNTALHYSVSHSNFEIVKLLLDADVNCVD
HQNKAGYTPIMLAALAAVEAEKDMRIVEELFGCGDVNAKASQAGQTALMLAVSHGRID
MVKGLLACGADVNIQDDEGSTALMCASEHGHVEIVKLLLAQPGCNGHLEDNDGSTALS
IALEAGHKDIAVLLYAHVNFKAQSPGTPRLGRKTS PGPTHRSFD"
```

ORIGIN

```
1 gcggccgaag aaaaacagga caaagaaacc tgtgtgaatt ttcagctttt caccttctct
61 gatttttattt cttcctcact cttcctttga cagtccctgc agtccgcctg agagagtaga
121 gaagaccccc tcccagaacc ttctgttaag gtctcccgcc tgacttctgt tagtggatgt
181 gactgtgtgc caggtgcctg ccgaagaccc ctgtcactga ctgtgccctt tgggaaagag
241 tcaacaatgg cctcctccta tggctgactt cgttcatgtt caggctctcc agcttgctag
301 tgagaattgc caaggattgg ttcatggcag gatagaacta aactgataga tgaaagctcc
361 acagtgtctc aagacggggt cccgttcatg ttaagcagtt ttccttctct caaaataaaa
421 aactagcagt ccttagggag gacagttttt tcttctcttt tctattctg gctccaacag
481 ctgtctcaac acagcccgag atggggagca gcctggcttc caccggcaat agctgtattg
541 tgggagtgtg aaagagaagc caccttttcc tccctctgcc caagccacct ggcccctttg
601 tcctttctcc tctcgtctct ctgaggttga atgcctttga gaacttgatg cataaaattt
661 gcatgactcc tcactccttt ctggatctcc cattggactc aagccagcat ggctcacacc
721 acaaaggtta acggcaagtgc ctcaggaaa gcaggtgata ttctcagtg agaccaggac
781 aaggaacaga aagaccctta ctttgtggag acccctatg gttatcaact agacttagat
841 ttctcaaat atgtggatga catacagaag ggaaatacca tcaaagact gaacatccag
901 aagaggcgga agccgtccgt gccatgccc gaaccagga ccacatctgg tcagcaaggt
961 atatggactt ccaactgaatc cctctcatcc tcaaacagt atgacaacaa gcagtgtccc
1021 aacttctca tagccagaag tcaagttaca tcaactcaa tctcaaagcc acctccccct
1081 ctggagacct cactcccttt tcttaccatc ccagaaaatc gacagctgcc acctccctca
1141 ccacaactcc caaagcataa ccttcatgtc accaagacac tgatggagac ccggagaaga
1201 ctggaacagg agagagccac catgcagatg acaccgggtg agttcagaag gccaggctg
1261 gccagttttg gaggcattgg caccacaagc tccctccctt cttttgtggg ttctggaaac
1321 cacaatcctg ccaagcacca gcttcagaat ggataccaag gtaatgggga ttatggtagc
1381 tatgccccag ctgctccccc cacttccctc atggggagct ccatccgcca cagccccctg
1441 agetcaggga tctccacccc agtgaccaac gtgagcccca tgcacctgca gcacatccgc
1501 gagcagatgg ccattgctct gaaacgcctg aaggagctgg aggagcaggt gcgaaccatc
1561 cctgtgctcc aggtaaagat ctctgtcttg caagaagaga aaaggcagtt ggtctcacag
1621 ctgaaaaacc aaagggtgc atcccagatc aatgtctgtg qtgtgaggga gcggtccat
1681 agtgccggga acgctcccca gctggacacg ctctccggg cccgaagaag tggcgggga
1741 ttatacatag actatgagga ggaagaaatg gagaccgtag aacagagcac gcagaggata
1801 aaggagtccc ggcaacttac agcagacatg caagccctgg agcagaagat ccaggacagc
1861 agctgtgagg cctcctcaga gctcaggagg aatggagagt gccggtctgt ggctgtgggt
```

```
1921 gccgaggaga acatgaacga catcgctcgtg taccacagag gctccaggtc ctgtaaggat
1981 gcagctgtag ggacacttgt tgagatgaga aattgtgggg tcagcgtgac agaggccatg
2041 cttggagtga tgactgaagc tgacaaagaa attgagctcc aacagcagac catagaagcc
2101 ttgaaggaaa agatctatcg cctagaagta cagcttagag aaaccaccca tgaccgggag
2161 atgactaaac tgaacaaga gctgcaggct gctggatcga ggaaaaaggt tgacaaagcc
2221 acgatggccc agccgcttgt tttcagttaag gtggtggagg cagtgggtga gaccagagac
2281 caaatggtcg gcagtcacat ggacctggtg gacacgtgtg ttgggacctc cgtggaaaca
2341 aacagtgtag gcctctcctg ccagcctgaa tgaagaata aagtcgtagg gcctgagctg
2401 cctatgaatt ggtggattgt taaggagagg gtggaatgc atgaccgatg tgctgggagg
2461 tctgtggaaa tgtgtgacaa gagtgtgagt gtggaagtca gcgtctgcaa aacaggcagc
2521 aacacagagg agtctgtgaa cgacctcaca ctctcaaga caaacttgaa tctcaaaaga
2581 gtgcggtcta tcggttgtgg agattgttct gttgacgtga ccgtctgctc tccaaaggag
2641 tgcgcctccc ggggcgtgaa cactgaggct gttagccagg tgaagctgc cgtcatggca
2701 gtgcctcgta ctgcagacca ggacactagg acagatttgg aacagggtga ccagttcacc
2761 aacaccgaga cggccaccct catagagtcc tgcaccaaca cttgtctaag cactttggac
2821 aagcagacca gcacccagac tgtggagacg cggacagtag ctgtaggaga aggcctgtc
2881 aaggacatca actcctccac caagacgcgg tcatttgggt ttggaacgtt gctttctggc
2941 cattctgggt ttgacaggcc atcagctgtg aagaccaaag agtcaggtgt ggggcagata
3001 aatattaacg acaactatct ggttgggtctc aaaatgagga ctatagcttg tgggccacca
3061 cagttgactg tggggctgac agccagcaga aggagcgtgg gggttgggga tgaccctgta
3121 ggggaatctc tggagaaccc ccagcctcaa gctccacttg gaatgatgac tggcctggat
3181 cactacattg agcgtatcca gaagctgtcg gcagaacagc agacactgct ggctgagAAC
3241 tacagtgaac tggcagaagc tttcggggaa cctcactcac agatgggctc cctcaactct
3301 cagctcatca gcacctgtc gtctatcaac tctgtcatga aatctgcaag cactgaagag
3361 ctgaggaaac ctgaactcca gaaaaccagt ctgggtaaaa tcacaggcaa ttatttggga
3421 tatacctgta agtgtggggg ccttcagtca ggaagtccct taagctcca gacatccag
3481 cctgagcaag aagtggggac ctcaagaagg aagccaatca gcagcctgga tgcttcccc
3541 actcaggaag gtacgtgtgc tccagtgaac ctgacagacg accagatcgc cgctggcctc
3601 tatgcatgta caaacaatga aagtacactg aagtccatca tgaagaagaa agatggtaac
3661 aaagattcaa atggcgcaaa aaagaatctt cagtttgttg gcattaatgg agggatgaa
3721 acaacttcaa gtgatgattc cagctcagat gaaagctctt cttccagatc agatgacgag
3781 tgtgatgtca ttgagtatcc tctgaagaa gaggaggagg aggaggatga agacactcgg
3841 ggaatgctga aagggcacca tgcagttaat attgaagggt tgaagtctgc cagggtggaa
3901 gatgaaatgc aggttcaaga atgtgaacct gagaaggtgg aaatcagaga gaggtatgaa
3961 ttaagtgaaa agatgttgtc tgcagtcaac ttactgaaaa atactataaa tgaccccaaa
4021 gctttgacca gcaaagatat gaggttctgt ctgaacaccc tccagcacga gtggttccgc
4081 gtgtccagtc agaagtcagc cattccagcc atggtggggg actacatagc tgcttttgag
4141 gtcatttccc cagatgtcct ccgctatgtc atcaacttgg cagacggcaa cggcactaca
4201 gccctccatt acagcgtgtc ccactccaac ttcgagattg tgaagctgct gttagatgcc
4261 gatgtgtgta atgtggatca ccagaacaag gcaggctaca ccccatcat gttggcggcc
4321 ctgcccgtcg tggaaagcaga gaaggacatg cggattgtgg aagaactctt tggctgtggg
4381 gatgtgaatg ccaaagctag tcaggcggga cagacggccc tcatgctggc ggtcagtcac
4441 ggacggatag acatggtgaa gggccttctg gcctgtgggg ctgatgtcaa catccaggat
4501 gacgagggct ccacggccct catgtgtgcc agcagcacg ggcacgtgga gattgtcaag
4561 ctgtgtctgg cccagcccg gtgcaacggt cacctagagg acaacgatgg cagcactgcg
4621 ctctcaatcg ccctggaagc aggacacaag gacatcgctg ttcttctgta tgcccatgtc
4681 aactttgcaa aagcccagtc tccgggcacc cctaggcttg gaaggaagac gtctcctggc
4741 cccacccacc gaggttcatt tgattgattg tatgcaata gccctttatt tacatgccac
4801 tattaagctg ctaattgttc ctgttggggt gacagatact gaatgtatac gtattgtgcc
4861 tgagctcacc agcaaacaga agcatcaagc ccagggttaa aggtgaagc tttcacagtg
4921 cagagactgc tagcctgggc acacgcacct cctttctggc cgtcttctgt gtagggcaca
4981 ctttaaccca gtctctgttg ctgttgagtc tctgtccgt tttgtacagt cacagggaaat
5041 tctgatctga aggggcacct tctgttcact ccacaaaagt ggtgtctggt tctcactgag
5101 acgttttaag atttttccac aaatatattat atgtactaaa tgtggaacca ttagaaagtt
5161 cttccaaaat ctcatccag catagttttg gatttttctt ttgtcttatt ttaaaataag
5221 gaagtcgaga tgactttgat catttgtaac ttgggccttg gccagacaaa gtataaaact
5281 tacaaaagaa tattctcatt tggctttaac taggtagatg taatatatga cttttataaa
5341 aaagggtatc tatatgaact tgacacagta ttttcagctt ttgtattcca tactaaagcc
5401 atgaagaact acacgtaaca tcatcatttg tattaattgc acaactccaa tgctaaaggt
5461 tggatttgtt tagaggaatc ggctctgtat ttgctctag agaaacacag tgttctcttt
5521 gtatttatgg attccttttt accgtgtcac atttactttg gtcctctatg tatttaaatg
5581 tttgaagtgc cttagactct tgccatattt tcaaaataaa attccattaa gctct
```

//

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record.**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☒ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.